

# WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5:	A 1	(11) International Publication Number:	WO 93/11773
A61K 31/57 // (A61K 31/57 A61K 31:165)	A1	(43) International Publication Date:	24 June 1993 (24.06.93)
<ul> <li>[21] International Application Number: PCT/EF</li> <li>[22] International Filing Date: 7 December 1992</li> <li>[30] Priority data: 91311761.0 18 December 1991 (18.12)</li> <li>[71] Applicant: AKTIEBOLAGET ASTRA [SE/SE]; Södertälje (SE).</li> <li>[72] Inventors: CARLING, Christer, Carl, Gustav; F 8, S-240 10 Dalby (SE). TROFAST, Jan, Will penkroken 34, S-226 47 Lund (SE).</li> <li>[74] Agents: HJERTMAN, Ivan et al.; AB Astra, P partment, S-151 85 Södertälje (SE).</li> </ul>	(07.12.9 2.91) ; S-151 Backväg liam ;	DE, DK, ES, FI, GB, HU, JP, MN, MW, NL, NO, NZ, PL, P European patent (AT, BE, CH GR, IE, IT, LU, MC, NL, PT BJ, CF, CG, CI, CM, GA, GN,  E  Published  With international search report.  Before the expiration of the time claims and to be republished in amendments.	KP, KR, LK, LU, MG, I, RO, RU, SD, SE, UA, DE, DK, ES, FR, GB, SE), OAPI patent (BF, ML, MR, SN, TD, TG).

#### (57) Abstract

Effective amounts of formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide are used in combination for simultaneous, sequential or separate administration by inhalation in the treatment of respiratory disorder.

## FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

ustria ustralia arbudos elgium urkina Faso ulgaria enin razil anada entral African Republic	FR GA GB GN GR HU IE IT JP KP	Gabon United Kingdom Guinea Greece Hungary Ireland Italy Japan Democratic People's Republic	MW NL NO NZ PL PT RO RU SD	Malawi Netherlands Norway New Zealand Poland Portugal Romania Russian Federation
arbados elgium urkina Faso ulgaria enin razil anada	GB GN GR HU IE IT JP	United Kingdom Guinea Greece Hungary Ireland Italy Japan	NO NZ PL PT RO RU	Norway New Zealand Poland Portugal Romania
elgium urkina Faso ulgaria enin razil anada	GN GR HU IE IT JP	Guinea Greece Hungary Ireland Italy Japan	NZ PL PT RO RU	New Zealand Poland Portugal Romania
urkina Faso ulgaria enin razil anada	GR HU IE IT JP	Greece Hungary Ireland Italy Japan	PL PT RO RU	Poland Portugal Romania
ulgaria enin enzil anada	HU IE IT JP	Hungary Ireland Italy Japan	PT RO RU	Portugal Romania
enin rozil anada	ie It Jp	treland Italy Japan	RO RU	Romania
razil anada	IT JP	Italy Japan	RO RU	Romania
anada	JP	Japan	RU	
				A Advention to A description of the second s
entral African Republic	KP	Democratic Puopic & Kupuniiu		Sudan
			SE	Sweden
ongo		of Korea		Slovak Republic
vitzerland	KR	Republic of Korea	SK	· •
Ste d'Ivoire	KZ	Kazaklistan	SN	Senegal
anicroon	LI	Liechtenstein	SU	Soviet Union
rechoslovakia -	LK	Srī Lanka	TD	Chad
zech Republic	រ.ប	Luxembourg	TG	Togo
<b>V</b>	MC	Monaco		Ukraine
<u>*</u>		Madagascar	US	United States of America
			٧N	Vict Nam
min.				
ei Ei	ch Republic many imark in	many MC mark MG in ML	many MC Monaco mark MG Madagascar in ML Mali	many MC Monaco UA mark MG Madagascar US

New combination of formoterol and budesonide.

5

•

¥.

4

#### Field of the invention

This invention relates to improvements in the treatment of mild as well as severe asthma and other respiratory disorders. More particularly, it relates to the use of a bronchodilator in combination with a steroidal anti-inflammatory drug for the treatment of respiratory disorders such as asthma, and to pharmaceutical compositions containing the two active ingredients. It emphasizes the use of a long-acting bronchodilator which provides rapid relief of symptoms.

#### Background of the invention

There have recently been significant advances in our 20 understanding of asthma. Despite many advances, both in awareness of the disease by doctors and patients alike, coupled with the introdction of very powerful and effective anti-asthma drugs, asthma remains a poorly understood and often poorly treated disease. Previously, 25 contraction of airway smooth muscles has been regarded as the most important feature of asthma. Recently there has been a marked change in the way asthma is managed, stemming from the fact that asthma is recognized as a chronic inflammatory disease. Uncontrolled airway 30 inflammation may lead to mucosal damage and structural changes giving irrversible narrowing of the airways and fibrosis of the lung tissue. Therapy should therefore be aimed at controlling symptoms so that normal life is possible and at the same time provide basis for treating

ŧ.

\*

the underlying inflammation.

The most common cause for poor control of asthma is poor compliance with the long-term management of chronic asthma, particularly with prophylatic treatments, such as inhaled steroids, which do not give immediate symptom relief. Patients will readily take  $\beta_2$ -agonist inhalers, since these provide rapid relief of symptoms, but often do not take prophylactic therapy, such as inhaled steroids, regularly because there is no immediate symptomatic benefit. They also counteract down regulation of  $\beta_2$ -adrenoceptor agonists.

Formoterol, (N-[2-hydroxy-5-[1-hydroxy-2-[[2-(4methoxyphenyl)-1-methylethyl]amino]ethyl]phenyl] 15 formamide), is an adrenoceptor agonist which selectively stimulates  $\beta_2$ -receptors, thus producing relaxation of bronchial smooth muscle, inhibition of the release of endogenous spasmogens, inhibition of oedema caused by endogenous mediators, and increased mucociliary 20 clearence. Inhaled formoterol fumarate acts rapidly, usually within minutes which gives the patient immediate confirmation that he has taken an adequat dose and thereby avoiding overdosing of both  $\beta$ -agonist and steroid. Inhaled formoterol also exerts a prolonged 25 bronchodilation, which in clinical trials has been demonstrated as up to 12 hours.

Budesonide, (16,17-butylidenebis(oxy)-11,21
dihydroxypregna-1,4-diene-3,20-dione), may be given in a high inhaled dose (up to 2 mg daily) with very low systemic effects, possibly because of its rapid metabolism. The high rapid systemic elimination of budesonide is due to extensive and rapid hepatic

metabolism. Long term clinical studies have shown that inhaled budesonide is a pharmacologically safe drug. High doses of inhaled budesonide are highly effective and well

PCT/EP92/02826

\*

tolerated when used in oral steroid replacement therapy. Budesonide represents a logical safe and effective therapy for long term control of asthma.

5 The inhaled route of administration enables the dose to be delivered directly to the airways. By this type of administration, it is possible to give a small dose and thereby minimizing unwanted side-effects. The drawbacks of the currently available bronchodilators are their relatively short duration of action. By using a compound with long duration e.g. formoterol it would be possible to avoid the nocturnal asthma, which so often causes considerable anxiety and debility to the patients. Formoterol gives less nocturnal waking than the commonly used short-acting agonists like salbutamol, terbutaline and the like. Formoterol has been registered for oral administration in Japan since 1986.

Pharmaceutical combinations of long-acting  $\beta_2$ -agonists and steroids are disclosed in two European applications, EP 416950 which discloses the combination of salmeterol and beclomethasone, and EP 416951 which discloses the combination of salmeterol and fluticasone propionate.

In Ann. Allergy 1989, 63 (3), p. 220-224 the use of a  $\beta_2$ -agonist, i.e. formoterol and a steroid, i.e. budesonide seperately are mentioned. It is not disclosed a pharmaceutical combination including both formoterol and budesonide, or the use of the two compounds in combination therapy. The use of a  $\beta_2$ -agonist and a steroid separately is also mentoined in Lung (1990), 168, no. supp, p. 105-110.

### Outline of the Invention

35

*¥* 

The present invention is based on the concept of a novel combination therapy whereby formoterol (and/or a

7

Ŧ,

25

35

physiologically acceptable salt and/or solvate thereof) and budesonide are administrated simultaneously, sequentially or seperately by inhalation. This combination has not only a greater efficiency and duration of bronchodilator action but the combination 5 also has a rapid onset of action. This new feature is of utmost importance in order to establish a higher compliance for patients and it provides a rescue medicine thereby avoiding the necessity for the patient of carrying two different inhalers. This simplifies life for 10 patients considerably and makes life more comfortable and secure. The rapid onset of the long-acting  $\beta_2$ -agonist gives the patient immediate confirmation that he has taken an adequate dose and thereby avoiding overdosing of both  $\beta_2$ -agonist and steroid. Since the use of formoterol 15 instead of salmoterol gives a much more rapid onset the combinations according to the invention have a number of advantages compared to the combinations disclosed i EP 416950 and EP 41651. The combination according to present invention permits a twice daily dosing regime as a basic 20 treatment of asthma, particularly nocturnal asthma.

The present invention provides a medicament containing, separately, or together, (i) formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and (ii) budesonide for simultaneous, sequential or separate administration by inhalation in the treatment of respiratory disorder.

The invention also provides a pharmaceutical composition for administration by inhalation in the treatment of respiratory disorder which composition comprises formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide.

According to another aspect of the invention there are provided pharmaceutical compositions comprising effective

5

amounts of formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide as a combined preparation for simultaneous, sequential or seperate administration by inhalation in the treatment of respiratory disorder.

3

¥

5

10

30

×

The invention further provides formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide for use in combination therapy by simultaneous, sequential or seperate administration by inhalation in the treatment of respiratory disorder.

Further the invention provides the use of formoterol (and/or a physiologically acceptable salt and/or solvate 15 thereof) in the manufacture of a medicament for combination therapy where formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide are administered simultaneously, sequentially or seperately by inhalation in the treatment 20 of respiratory disorder and the use of budesonide in the manufacture of a medicament for combination therapy where formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide are administered simultaneously, sequentially or separately by inhalation in the treatment of respiratory disorder. 25

The invention additionally relates to the use of formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide in the manufacture of a medicament for combination therapy for simultaneous, sequential or seperate administration of formoterol and budesonide by inhalation in the treatment of respiratory disorder.

According to a further feature of the invention there is provided a method of treating respiratory disorder which comprises the simultaneous, sequential or separate

•

3

20

25

30

35

administration by inhalation of effective amounts of formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide.

Suitable physiologically salts of formoterol include acid addition salts derived from inorganic and organic acids, such as the hydrochloride, hydrobromide, sulphate, phosphate, maleate, fumarate, tartrate, citrate, benzoate, 4-methoxybenzoate, 2- or 410 hydroxybenzoate, 4-chlorobenzoate, p-toluenesulphonate, methanesulphonate, ascorbate, salicylate, acetate succinate, lactate, glutarate, gluconate, tricarballylate, hydroxynaphthalenecarboxylate or oleate. Formoterol is preferably used in the form of its fumarate salt and as a dihydrate.

The ratio of formoterol to budesonide used according to the invention is preferably within the range of 1:4 to 1:70. The two drugs may be administered separately in the same ratio.

The intended dose regimen is a twice daily administration, where the suitable daily dose of formoterol is in the range of 6 to 100 µg with a preferred dose of 6-48 µg and the suitable daily dose for budesonide is 50 to 4800 µg with a preferred dose of 100-1600 µg. The particular dose used will strongly depend on the patient (age, weight etc) and the severity of the disease (mild, moderate, severe asthma etc).

For administration, the combination is suitably inhaled from a nebulizer, from a pressurized metered dose inhaler or as a dry powder from a dry powder inhaler (e.g. as sold under the trade mark Turbuhaler) or from a dry powder inhaler utilizing gelatine, plastic or other capsules, cartridges or blister packs.

A diluent or carrier, generally non-toxic and chemically inert to the medicament e.g. lactose, dextran, mannitol or glucose or any additives that will give the medicament a desired taste, can be added to the powdered medicament.

5

10

'n

4

Examples of the preparation of suitable dosage forms according to the invention include the following: Formoterol fumarate dihydrate and budesonide (optionally premicronized) are mixed in the proportions given above. The agglomerated, free-flowing micronized mixture may be filled into dry powder inhaler such as sold under the trade mark Turbuhaler. When a capsule system issued, it is desirable to include a filler in the mixture.

- The micronized mixture may be suspended or dissolved in a 15 liquid propellant mixture which is kept in a container that is sealed with a metering valve and fitted into a plastic actuator. The propellants used may be chlorofluorocarbons of different chemical formulae. The most frequently used chlorofluorocarbon propellants are 20 trichloromonofluoromethane (propellant 11), dichlorodifluoromethane (propellant 12), dichlorotetrafluoroethane (propellant 114), tetrafluoroethane (propellant 134a) and 1,1-difuoroethane (propellant 152a). Low concentrations of a 25 surfactant such as sorbitan trioleate, lecithin, disodium dioctylsulphosuccinate or oleic acid may also be used to improve the physical stability.
- The invention is further illustrated by way of example with reference to the following Examples.

#### Example 1 - Dry Powder Inhaler (Turbuhaler)

35	Active ingredient	<u>Per dose</u>
•	Formoterol (as fumarate dihydrate)	12 μg
•	Budesonide	200 μg

The storage unit of the inhaler is filled with sufficient for at least 200 doses.

5	Active ingredient	Per dose
5	Formoterol (as fumarate dihydrate)	24 µg
	Budesonide	200 µg
	The storage unit is filled with sufficient	for at least
	200 doses.	
10	200 00000	•
10	Active ingredient	<u>Per dose</u>
	Formoterol (as fumarate dihydrate)	12 μg
	Budesonide	100 µg
	The storage unit is filled with sufficient	for at least
15	200 doses.	
10	200 40505.	
	Example 2 - Metered dose inhaler	•
	Active ingredient	Per dose
20	Formoterol (as fumarate dihydrate)	12 μg
20	Budesonide	200 <b>µ</b> g
	Stabilizer	0.1 - 0.7  mg
	Propellant	$25 - 100 \mu l$
25		
	Active ingredient	Per dose
	Formoterol (as fumarate dihydrate)	24 µg
	Budesonide	200 <b>µ</b> g
	Stabilizer	0.1 - 0.7  mg
30	Propellant	$25 - 100 \mu l$
	Active ingredient	<u>Per dose</u>
	Formoterol (as fumarate dihydrate)	12 μg
35	Budesonide	200 µg
	Stabilizer	0.1 - 0.7  mg
	Propellant	$25 - 100 \mu l$

PCT/EP92/02826

## Example 3 - Metered dose dry powder formulation

	Active ingredient	Per dose
	Formoterol (as fumarate dihydrate)	12 μg
5	Budesonide	200 <b>µ</b> g
	Lactose	up to 5, 12.5 or 25 mg
	Active ingredient	<u>Per dose</u>
10	Formoterol (as fumarate dihydrate)	24 <b>µ</b> g
	Budesonide	200 μg
	Lactose	up to 5, 12.5 or 25 mg
15	Active ingredient	Per dose
	Formoterol (as fumarate dihydrate)	$12 \mu g$
	Budesonide	100 μg
	Lactose	up to 5, 12.5 or 25 mg

2.0

WO 93/11773

10

#### CLAIMS

- 1. A medicament containing, seperately or together, (i) formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and (ii) budesonide for simultaneous, sequential or seperate administration by inhalation in the treatment of respiratory disorder.
- 2. A pharmaceutical composition for administration by inhalation in the treatment of respiratory disorder which composition comprises formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide.
- 3. A pharmaceutical composition comprising effective amounts of formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide as a combined preparation for simultaneous, sequential or separate administration by inhalation in the treatment of respiratory disorder.
  - 4. Formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide for use in combination therapy by simultaneous, sequential or separate administration by inhalation in the treatment of respiratory disorder.

25

35

- 5. The use of formoterol (and/or a physiologically acceptable salt thereof) in the manufacture of a medicament for combination therapy where formoterol (and/or physiologically acceptable salt and/or solvate thereof) and budesonide are administered simultaneously, sequentially or separately by inhalation in the treatment of respiratory disorder.
  - 6. The use of budesonide in the manufacture of a medicament for combination therapy where formoterol

(and/or a physiologically acceptable salt and/or solvate thereof) and budesonide are administered simultaneously, sequentially or separately by inhalation in the treatment of respiratory disorder.

11

5

10

\*

7. The use of formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide in the manufacture of a medicament for combination therapy for simultaneous, sequential or separate administration of formoterol and budesonide by inhalation in the treatment of respiratory disorder.

		INTERNATIONA	L SEARCH REPORT International Application No	PCT/EP 92/02826
I. CLASSII	CATION OF SUB	JECT MATTER (if several classificati	ion symbols apply, indicate all) <sup>6</sup>	
_		ent Classification (IPC) or to both Nation 57; //(A61K31/		
II. FIELDS	SEARCHED			
<del></del>		Minimum Do	cumentation Searched?	
Classification	on System	T	Classification Symbols	· · · · · · · · · · · · · · · · · · ·
Int.C1.	5	A61K		
•			ther than Minimum Documentation ents are Included in the Fields Searched <sup>8</sup>	
m. Docum	IENTS CONSIDER	ED TO BE RELEVANT <sup>9</sup>		
Category o	Citation of I	Document, $^{11}$ with indication, where appr	ropriate, of the relevant passages 12	Relevant to Claim No.13
X	ANNALS vol. 63 pages 2 JEAN H. agonist treatme cited i see pag see pag * page  LUNG (U vol. 16 pages 1 NILS SV beta-ag	OF ALLERGY 3, no. 3, September 19 220 - 224 . MARSAC ET AL. 'Inhalts and inhaled steroid ent of asthma' in the application ge 221, column 2; tablue 221, column 3, line 223; summary *  USA) SS, no. SUPP, 1990, NE. 195 - 110 /EDMYR 'The current plants in the management of the application of the application.	189, led beta is in the le 1 le 1 - line 15  W YORK lace of	1-7
"A" documents consinue consinu	idered to be of particler document but puby date ment which may through its cited to establish on or other special rement referring to an means ment published prior than the priority date.	eneral state of the art which is not cular relevance lished on or after the international ow doubts on priority claim(s) or a the publication date of another eason (as specified) oral disclosure, use, exhibition or to the international filing date but te claimed	"I" later document published after the in or priority date and not in conflict we cited to understand the principle or to invention  "X" document of particular relevance; the cannot be considered novel or cannot involve an inventive step  "Y" document of particular relevance; the cannot be considered to involve an indocument is combined with one or meants, such combination being obvious in the art.  "&" document member of the same patent.	theory underlying the claimed invention t be considered to claimed invention nventive step when the nore other such docu- ous to a person skilled at family
Date of the Ac	•	the International Search	Date of Mailing of this International	Search Report
07 APRIL 1993			1281414	

Signature of Authorized Officer

LEHERTE C.F.M.

Form PCT/ISA/210 (second sheet) (Jamesry 1985)

EUROPEAN PATENT OFFICE

International Searching Authority



# SECOND CORRECTED VERSION \*\*

## **PCT**

# WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5:
A61K 31/57 // (A61K 31/57
A61K 31:165)

A1
(11) International Publication Number: WO 93/11773
(43) International Publication Date: 24 June 1993 (24.06.93)
(21) International Application Number: PCT/EP92/02826 (81) Designated States: AT, AU, BB, BG, BR, CA, CH, CS,

(22) International Filing Date: 7 December 1992 (07.12.92)

was filed:

(30) Priority data:
91311761.0
18 December 1991 (18.12.91)
EP
(34) Countries for which the regional
or international application

SE et al.

'(71) Applicant: AKTIEBOLAGET ASTRA [SE/SE]; S-151 85 Södertälje (SE).

(72) Inventors: CARLING, Christer, Carl, Gustav; Backvägen 8, S-240 10 Dalby (SE). TROFAST, Jan, William; Vapenkroken 34, S-226 47 Lund (SE).

(74) Agents: HJERTMAN, Ivan et al.; AB Astra, Patent Department, S-151 85 Södertälje (SE).

(81) Designated States: AT, AU, BB, BG, BR, CA, CH, CS, DE, DK, ES, FI, GB, HU, JP, KP, KR, LK, LU, MG, MN, MW, NL, NO, NZ, PL, RO, RU, SD, SE, UA, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, SN, TD, TG).

**Published** 

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: NEW COMBINATION OF FORMOTEROL AND BUDESONIDE

(57) Abstract

Effective amounts of formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide are used in combination for simultaneous, sequential or separate administration by inhalation in the treatment of respiratory disorder.

## FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinca	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JР	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU -	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic	SD	Sudan
CG	Congo		of Korca	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SI	Slovenia
Cl	Côte d'Ivoire	KZ	Kazakhstan	SK	Slovakia
CM	Cameroon	LI	Liechtenstein	SN	Senegal
CN	China	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
CZ	Czech Republic	LV	Latvia	TJ	Tajikistan
DE	Germany '	MC	Monaco	TT	Trinidad and Tobago
DK	Denmark	MD	Republic of Moldova	UA	Ukraine
ES	Spain	MG	Madagascar	US	United States of America
FI	Finland	, ML	Mali	UZ	Uzbekistan
FR	France	MN	Mongolia	VN	Vict Nam
GA	Gabon		-		